

# Aura MLS Radiance Average Retrievals (RAR) HO<sub>2</sub> product guideline

## Summary

Useful vertical range: 10 – 0.0032 hPa (daytime)  
day-minus-night differences required between 10 and 1 hPa

1–0.0032 hPa (nighttime)

Latitude coverage: 85°S to 85°N

Vertical resolution: varying from ~4 km to 14 km

Significant averaging (monthly means) is required to obtain scientifically useful data.

Contact: Luis Millán

Jet Propulsion Laboratory, California Institute of Technology

[luis.f.millan@jpl.nasa.gov](mailto:luis.f.millan@jpl.nasa.gov)

## Introduction

A description of the retrieval methodology is given by:

Millán et al. (2015), Stratospheric and mesospheric HO<sub>2</sub> observations from the Aura Microwave Limb Sounder, *Atmos. Chem. Phys.*, 15, 2889–2902, 2015, doi:10.5194/acp-15-2889-2015

In short, the retrieval algorithm produces a pair of zonal-mean abundance fields for each day, one for the daytime part of the orbit and the other for the nighttime part on a grid with 6 surfaces per decade change in pressure (~3 km). These are obtained from a 10° latitude bin zonal mean of radiances (sorted by solar zenith angle) interpolated onto a 6 surface per decade pressure grid using the limb tangent pressure from the standard production data.

To minimize biases, day-minus-night differences between 10 and 1 hPa must be used as a more accurate measure of daytime HO<sub>2</sub>.

Due to the small spectral signature of HO<sub>2</sub> in the MLS radiances, significant averaging (such as monthly zonal means) is required to obtain scientifically useful results.

## Precision, Accuracy, and Vertical Resolution

In the usable pressure range, the vertical resolution varies from about 4 km at 10 hPa to about 14 km at 0.0032 hPa. Daily precision for a 10° latitude bin ranges from 0.1 ppbv in the upper stratosphere to up to 8 ppbv in the upper mesosphere, dropping to ~1.4 ppbv and ~0.5 ppbv for monthly and yearly averages,

respectively. Between 10 and 0.1 hPa, for both daytime and nighttime cases, the total systematic error is around 0.04 ppbv (up to  $\sim 10 \times 10^6 \text{ molec cm}^{-3}$ ), while for lower pressures (i.e., higher altitudes) the systematic error is as large as 1.2 ppbv ( $\sim 0.2 \times 10^6 \text{ molec cm}^{-3}$ ).

## Data Format

All the MLS HO<sub>2</sub> data described here can be found at the NASA Goddard Space Flight Center Earth Sciences (GES) Data and Information Services Center DISC [website](#).

All the data described here are stored in netCDF files.

The data are stored in files named according to the convention

MLS-Aura.L3ZMRAR-HO<sub>2</sub>-v06-VV>-c<CC>.<yyyy>d<ddd>.nc4

where L3ZMRAR stands for Level 3 Zonal Means Radiance Average Retrievals, v06-VV>-c<CC> is the version and cycle number. The files are produced on a one-day granularity and named according to the observation date, where <yyyy> is the four-digit calendar year and <ddd> is the day number in that year (001 = 1 January).

Each file contains two swaths: Daytime and Nighttime. Each swath contains the following fields:

|                    |                                |   |
|--------------------|--------------------------------|---|
| Average            | retrieved HO <sub>2</sub> data | [vmr]   |
| Average ND         | retrieved HO <sub>2</sub> data | [10 <sup>6</sup> molecules cm <sup>-3</sup> ] |
| Error              | precision                      | [vmr]   |
| Error ND           | precision                      | [10 <sup>6</sup> molecules cm <sup>-3</sup> ] |
| lat                | latitude                       | [-85°, -75°, -65°, ... , 85°]                 |
| lev                | pressure levels                | [hPa]   |
| Solar_Zenith_Angle | solar zenith angle             | [deg]   |
| Local_Solar_Time   | local solar time               | [hours]                                       |

## Data Screening

Bad data were set to -999.99 and should be avoided.

Due to the small signal-to-noise ratio many negative values are found throughout this data set. These values need to be included in any scientific study to avoid high biases in averages derived from these data.

©2026 California Institute of Technology. Government sponsorship acknowledged.

The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018D0004).